

INFORMATION SHEET

Home Care Package Providers

Towards the end of life, palliative care patients with Level 3 & 4 Home Care Packages may need subcutaneous medicines for breakthrough symptom control. Carers (unpaid), when adequately trained, can assist in this role.

caring@home resources, targeted to health care professionals, community service providers and carers aim to support carers to help safely manage breakthrough symptoms that require subcutaneous medicines for home-based patients.

The resources are free, evidence-based and facilitate high-value care; they support person-centred outcomes and improve service performance efficiency and effectiveness. Using these resources to support home-based consumers and their carers helps providers to meet Aged Care Quality Standards #1,2,3,7 & 8 and the Charter of Aged Care Rights.

Home Care Package providers can use the *caring@home* resources to educate carers and support high-quality palliative care practice.

The resources are applicable in all jurisdictions across Australia.

Why teach carers to manage subcutaneous medicines?

One of the most frequent reasons that community-based palliative patients are transferred to inpatient units in the last week of life is because their symptoms cannot be adequately controlled at home. [1]

Research confirms that with standardised support from health care professionals and adequate information and resources, carers can be taught the skills to help manage breakthrough symptoms safely for the person for whom they are caring. [2,3]

Importantly, this can improve outcomes for patients and carers report feeling a great deal of satisfaction from contributing to symptom control, impacting positively on their bereavement. [4]

Resources to support Home Care Package providers

- *Guidelines for the handling of palliative care medicines in community services*
- Example policy and procedures: *Supporting carers to help manage breakthrough symptoms safely using subcutaneous medicines in the home*
- Online education modules for nurses
- A range of resources that can be used by nurses to train carers to manage breakthrough symptoms safely using subcutaneous medicines.

caring@home aims to improve the quality of palliative care service delivery across Australia.

The national project has developed standardised and free resources that support people to be cared for and to die at home, if that is their choice.



"caring@home resources can be used to improve outcomes for patients and outcomes for their carers in the grieving process. These are good resources to enable the patient to be cared for and die at home if they wish."

A GP who has used the *caring@home* resources

For more information:

T: 1300 600 007

caringathome@health.qld.gov.au

caringathomeproject.com.au

caring@home resources

Community service providers	<i>Guidelines for the handling of palliative care medicines in community services</i>	These guidelines, developed by NPS MedicineWise for <i>caring@home</i> , can be used by community service providers to inform the development of detailed protocols and procedures tailored to the requirements of individual services.
	Example policy and procedures: <i>Supporting carers to help manage breakthrough symptoms safely using subcutaneous medicines in the home</i>	This document may be used by community service providers to develop and/or review relevant documentation within their own organisation's policy and procedure framework.
Health care professionals	Online education modules	The online education modules educate nurses to train carers to manage breakthrough symptoms safely using subcutaneous medicines.
	Palliative care symptom management medicines for Australians living in the community	A consensus-based list of medicines suitable for use in the community for the management of terminal symptoms, developed by ANZSPM.
	pallIMEDS	This free app, based on the ANZSPM medicine list, and developed by NPS MedicineWise familiarises primary care prescribers and community pharmacists with eight palliative care medicines used for management of terminal symptoms.
Carers	A practical handbook for carers: <i>Helping to manage breakthrough symptoms safely using subcutaneous medicines</i>	The handbook provides written and pictorial material with all the information a carer needs to help manage breakthrough symptoms safely using subcutaneous medicines.
	Writing a label, opening an ampoule and drawing up medicine: <i>A step-by-step guide</i>	This illustrated guide explains how to label a syringe correctly, open an ampoule and draw up medicine using a step-by-step approach.
	Giving medicine using a subcutaneous cannula: <i>A step-by-step guide</i>	This illustrated guide explains how to give medicine through a subcutaneous cannula using a step-by-step approach.
	Medicines diary	The medicines diary is for carers to record all the subcutaneous medicines that are given.
	Colour-coded labelling system	The colour-coded labelling system acts as a safety check to ensure the correct medicine is given for each breakthrough symptom. It includes sticky labels for syringes and the Symptoms and medicines: <i>Colour-coded fridge chart</i> .
	A practice demonstration kit	The demonstration kit is used to practise giving medicines through a subcutaneous cannula.
	Short training videos	The videos show how to do each step when giving subcutaneous medicine.
	Translated resources	Select carer resources are available online in the following languages: Arabic, Greek, Italian, Simplified Chinese, Traditional Chinese, Tagalog and Vietnamese.

1. Rosenberg JP, Bullen T, Maher K. Supporting family caregivers with palliative symptom management: A qualitative analysis of the provision of an emergency medication kit in the home setting. *Am J Hosp Palliat Care*. 2015; 32(5): 484-9.
2. Healy S, Israel F, Charles M, Reymond L. Laycarers can confidently prepare and administer subcutaneous injections for palliative care patients at home: A randomized controlled trial. *Palliat Med*. 2018; 32(7):1-8.
3. National Institute for Health Research, Journals Library. CARer-Administration of as-needed sub-cutaneous medication for breakthrough symptoms in home-based dying patients: a UK study (CARIAD). Available from: <https://www.journalslibrary.nihr.ac.uk/programmes/hta/151037/#/> [Accessed 6 February 2017].
4. Healy S, Israel F, Charles MA, Reymond L. An educational package that supports laycarers to safely manage breakthrough subcutaneous injections for home-based palliative care patients: development and evaluation of a service quality improvement. *Palliat Med*. 2013; 27(6): 562-570.