

Medicines diary 用药日记





Copyright

This document is licensed under a Creative Commons Attribution-NonCommercial-NoDerivs 3.0 Australia licence. To view a copy of this licence visit:

https://creativecommons.org/licenses/by-nc-nd/3.0/au/

© Brisbane South Palliative Care Collaborative 2018

In essence you are free to copy, distribute and transmit the work in its current form for non-commercial purposes. You must attribute the work in the manner specified by the authors. You may not alter, transform, or build on this work.

Acknowledgements

This diary was developed as part of *caring@home*, a project undertaken by a consortium of Metro South Health, operating through Brisbane South Palliative Care Collaborative (lead agency), Aged & Community Services Australia, Australian Primary Health Care Nurses Association, Flinders University (through CareSearch), Leading Age Services Australia, National Prescribing Service, Pharmaceutical Society of Australia, The Royal Australian College of General Practitioners and University of Technology Sydney.

We acknowledge and thank all the health professionals and carers who contributed to the development of this diary.

Funding statement

caring@home is funded by the Australian Government, Department of Health.

Suggested reference

Brisbane South Palliative Care Collaborative. *Medicines diary*. Brisbane: Brisbane South Palliative Care Collaborative; 2018.

Enquiries

All enquiries about this document should be directed to:

Brisbane South Palliative Care Collaborative

T: 1300 600 007

E: caringathome@health.qld.gov.au

Disclaime

This diary has been developed specifically for use to assist carers with recording subcutaneous medicines given to the person they are caring for.

While the Brisbane South Palliative Care Collaborative has exercised due care in ensuring an accurate template for recording subcutaneous medicines in the diary, the information in the diary is a general guide only to appropriate practice, to be followed subject to the clinician's judgement and the carer's preference in each individual case.

The Brisbane South Palliative Care Collaborative does not accept any liability for any injury, loss, or damage incurred by use of, or reliance upon, the information provided within this diary.



版权

本文件采用知识共享署名 - 非商业性 - NoDerivs 3.0澳大利亚许可协议授权。如欲查看此许可证,请访问:

https://creativecommons.org/licenses/by-nc-nd/3.0/au/

© Brisbane South Palliative Care Collaborative 2018

大体上,您可以出于非商业目的自由复制、分发和传输当前形式的作品。您必须以作者指定的方式指明作品的归属。您不得改变、转换或基于本作品进行创作。

鸣谢

本日记是作为caring@home的一部分编写的,这是由Metro South Health联盟开展的一个项目,通过以下机构执行: 布里斯班南部姑息治疗协作机构(Brisbane South Palliative Care Collaborative) (牵头机构)、澳大利亚老龄和社区服务(Aged & Community Services Australia)、澳大利亚初级保健护理护士协会(Australian Primary Health Care Nurses Association)、弗林德斯大学(Flinders University)(通过CareSearch)、澳大利亚领先老龄服务(Leading Age Services Australia)、全国处方服务署(National Prescribing Service)、澳大利亚药学会(Pharmaceutical Society of Australia)、澳大利亚皇家全科医学院(The Royal Australian College of General Practitioners)和悉尼科技大学(University of Technology Sydney)。

我们感谢所有为本日记的编写做出贡献的健康专业人士和照顾者。

经费声明

caring@home 由澳大利亚政府卫生部提供经费。

建议参考资料

Brisbane South Palliative Care Collaborative。《用药日记》。布里斯班: Brisbane South Palliative Care Collaborative; 2018。

查询

有关本文件的所有问题,请联系:

Brisbane South Palliative Care Collaborative

电话: 1300 600 007

电邮: caringathome@health.qld.gov.au

免责声明

创建本日记之目的是专门为照顾者们使用,协助他们记录用於被照顾的人的皮下注射药物。

虽然Brisbane South Palliative Care Collaborative在确保日记在提供记录皮下注射药物的准确模板方面已经尽责尽力,但本日记只是适当实践的一般指南,在各病例中应该按临床医生的判断和照顾者的优先程序实施。

对于因使用或依赖本日记中提供的信息而导致的任何伤害、损失或损害, Brisbane South Palliative Care Collaborative不承担任何责任。

Introduction




This diary is one part of the *caring@home* package for carers.

It is used to record all subcutaneous medicines given to a person to help manage their breakthrough symptoms.

The diary:

- Allows you to keep track of all subcutaneous medicines used and how well they worked
- Helps the health care team to assess if medicines were effective and if medicines need to be changed
- Enables communications between you, the person you are caring for and the health care team about the medicines used

It is important that the diary is completed each time you give a medicine for a breakthrough symptom.



引言



本日记是给照顾者的 *caring@home* 成套教材的一部分。 它用于记录给予被照顾的人的所有皮下注射药物,以帮助管控他们的突破性症状。

日记:

- •让您可以记录使用的所有皮下注射药物及其疗效
- •帮助医疗团队评估药物是否有效,以及是否需要更换药物
- 协助您、您所照顾的人和医疗团队之间在所用药物方面的沟通

重要的是,每次因突破性症状给药时,您都要在日记上记录下来。





Colour-coded labelling system


To ensure that you have selected the right medicine for a particular breakthrough symptom, always read the label on the syringe.

The colour-coded labelling system, shown below, acts as an extra safety check. It helps you to select the correct medicine for each breakthrough symptom. It includes:

Colour-coded sticky labels for filled syringes



• Symptoms and medicines: Colour-coded fridge chart





彩色编码标签系统

请务必阅读注射筒上的标签,以确保您对特定突破性症状选择了正确的药物。

如下所示,彩色编码标签系统可作为额外的安全检查。它可以帮助您为每个突破性症状选择正确的药物。这包括:

•贴在装药注射筒上的彩色编码标签



• 症状和药物: 彩色编码冰箱图表

caring @home 症状和药物:彩色编码冰箱图表 药物 联系方式: Midozalam 症状 症状 症状 症状 焦虑 24小时电话号码: 药物 药物 氯化钠 0.9% Symptom / 症状 Pain – 疼痛 Shortness of breath - 呼吸急促 症状 症状 Noisy'rattly' breathing - 呼吸"咯咯"作响 冲洗注射 Nausea and/or vomiting – 恶心和 / 或呕吐 Restlessness/agitation – 烦躁 / 躁动不安 Anxiety – 焦虑 有白色背景的黑文本 浅橙鲑鱼色 Confusion – 思维混乱


caring@home由澳大利亚政府卫生部提供经费并由Brisbane South Palliative Care Collaborative领导。

本资讯改编自澳大利亚卫生保健安全和质量委员会(Australian Commission on Safety and Quality in Health Care)之《2015年可注射药物,液体和管线的用户应用标签的国家标准》(National Standard for User-applied Labelling of Injectable Medicines, Fluids and Lines. 2015)。

How to complete this diary




This example shows you how to complete the diary for each medicine given.



如何填写本日记



这个例子向您展示了如何为所给的每种药物填写日记。



评论/您想要注意或提及的其他事项	
(2前, (0-10) "约20分钟后" 症状评级 (0-10) (0-10) (0-10) 1 2 3 4 5 6 7 8 9 10 0 = 没有症状痛苦 10 = 可能是最糟糕的症状痛苦	
(0-10) (0-10) (0-10) 0 1 2 3 4 0 - 1 2 3 4 (0-∞有)	
给药原因 •疼痛 •呼吸急促 •呼吸"咯咯"作响 •恶心和/或呕吐 •烦躁/躁动不安 •焦虑 •思维混乱或新的意识混乱 •其他	
震	
	
时间	
	

评论/您想要注意或提及的其他事项			
(2前) (920分钟后) (0-10) (0-10) (0-10) (0-10) 1 2 3 4 5 6 7 8 9 10 0 = 没有症状痛苦 10 = 可能是最糟糕的症状痛苦			
/之門, (0-10) (0-10) 0 1 2 3 4 0 = 沒有			
给药原因 •疼痛 •呼吸急促 •呼吸"咯咯"作响 •恶心和/或呕吐 •烦躁/躁动不安 •焦虑 •思维混乱或新的意识混乱			
影			
松			
回回			
田田			

评论/您想要注意或提及的其他事项	
(2前, 10) (920分钟后, 12, 14, 14, 15 (0-10) (0-10) (0-10) (0-10) 1 2 3 4 5 6 7 8 9 10 10 = 28 4 5 6 7 8 9 10	
(0-10) (0-10) (0-10) (0-1 2 3 4 (0-1 2 3 4 (0-2) (0-2) (0-2)	
给药原因 •疼痛 •呼吸急促 •呼吸"路路"作响 •恶心和/或呕吐 •烦躁/躁动不安 •焦虑 •思维混乱或新的意识混乱 •其他	
展	
透	
回	
田田	

评论/您想要注意或提及的其他事项			
(2前) (920分钟后) (0-10) (0-10) (0-10) (0-10) 1 2 3 4 5 6 7 8 9 10 0 = 没有症状痛苦 10 = 可能是最糟糕的症状痛苦			
/之門, (0-10) (0-10) 0 1 2 3 4 0 = 沒有			
给药原因 •疼痛 •呼吸急促 •呼吸"咯咯"作响 •恶心和/或呕吐 •烦躁/躁动不安 •焦虑 •思维混乱或新的意识混乱			
影			
松			
回回			
田田			



Notes
备注